

Register Number:

Date:20-11-2020

ST. JOSEPH'S COLLEGE (AUTONOMOUS), BANGALORE - 27

B.Sc. STATISTICS – III SEMESTER SEMESTER EXAMINATION – NOVEMBER 2020

ST 318 - STATISTICAL INFERENCE - I

Time: 21/2hrs

Max:70 Marks

This question paper has TWO printed pages and THREE parts

SECTION - A

I Answer any FIVE of the following:

 $5 \times 3 = 15$

- 1. Differentiate between statistic and parameter. Give an example for each.
- 2. What do you mean by invariance property of an estimator? Mention its application.
- 3. Define relative efficiency of an estimator. Mention its range.
- 4. Define likelihood function and obtain an expression for log-likelihood function for geometric distribution
- 5. Define Confidence Interval and Confidence Coefficient
- 6. Define size of a test. How level of significance is different from size of the test?
- 7. What is p-value? How is it useful?

SECTION - B

II Answer any FIVE of the following:

 $5 \times 7 = 35$

- 8. A) Examine whether sample variance $s^2 = \frac{1}{n} \sum_{i=1}^{n} (x_i \overline{x})^2$ is an unbiased estimator of population variance σ^2 , when a sample is drawn from N (μ , σ^2).
 - B) What are the criteria of a good estimator?

(5+2)

- 9. A) Obtain the relationship between mean square error and variance of an estimator
 - B) Let $x_1, x_2, ..., x_n$ be the random sample of size n from Bernoulli distribution with parameter p. Examine whether \bar{X} is a consistent estimator for p or not. (3+4)
- 10. A) Prove that Binomial distribution belongs to one parameter exponential family.
 - B) Distinguish between simple hypothesis and composite hypothesis.

(5+2)

- 11. A) Explain the procedure to obtain the maximum likelihood estimator.
 - B) Derive moment estimators of a and b for U(a, b).

(4+3)

	12. A) Explain the Pivot method for finding the confidence interval.	
	B) What happens to the confidence interval when confidence level $lpha$ increases	? (5+2)
	13. A) Write down the confidence interval for proportion and explain the notations	
	B) Derive 100(1 - α) % level of significance confidence interval for population n	ean when
	σ is unknown	(2+5)
	14. A) Explain the types of errors involved in testing of hypothesis problem with ar	example.
	Deduce relation between type II error and power	
	B) Define test function	(5+2)
	SECTION - C	
111	Answer any TWO of the following: 2 x	10 = 20
	Answer any TWO of the following: 2 x 15. A) Let $x_1, x_2,, x_n$ be the random sample of size n from Poisson with par	
111	15. A) Let $x_1, x_2,, x_n$ be the random sample of size n from Poisson with par Examine whether \bar{X} is a MVUE or not.	ameter θ. (7)
111	 15. A) Let x₁, x₂,, x_n be the random sample of size n from Poisson with par Examine whether X̄ is a MVUE or not. B) Write down formula for 100(1-α) % confidence interval for the difference being the confidence interval for the confidence interval for the difference being the confidence interval for the difference being the confidence interval for th	ameter θ. (7)
	15. A) Let $x_1, x_2,, x_n$ be the random sample of size n from Poisson with par Examine whether \bar{X} is a MVUE or not.	ameter θ. (7)
111	 15. A) Let x₁, x₂,, x_n be the random sample of size n from Poisson with par Examine whether X̄ is a MVUE or not. B) Write down formula for 100(1-α) % confidence interval for the difference being the confidence interval for the confidence interval for the difference being the confidence interval for the difference being the confidence interval for th	ameter θ. (7) ween two (3)
111	 15. A) Let x₁, x₂,, x_n be the random sample of size <i>n</i> from Poisson with par Examine whether X̄ is a MVUE or not. B) Write down formula for 100(1-α) % confidence interval for the difference being population means (Consider independent case only) 	ameter θ. (7) ween two (3)
111	 15. A) Let x₁, x₂,, x_n be the random sample of size <i>n</i> from Poisson with par Examine whether X̄ is a MVUE or not. B) Write down formula for 100(1-α) % confidence interval for the difference being population means (Consider independent case only) 16. A) Suppose that x₁, x₂,, x_n from a random sample from a geometric distribution. 	ameter θ. (7) ween two (3) ution with
III	 15. A) Let x₁, x₂,, x_n be the random sample of size <i>n</i> from Poisson with par Examine whether X̄ is a MVUE or not. B) Write down formula for 100(1-α) % confidence interval for the difference being population means (Consider independent case only) 16. A) Suppose that x₁, x₂,, x_n from a random sample from a geometric distribution parameter θ. Derive maximum likelihood estimator of θ. 	ameter θ. (7) ween two (3) ution with (5)
III	 15. A) Let x₁, x₂,, x_n be the random sample of size <i>n</i> from Poisson with par Examine whether X̄ is a MVUE or not. B) Write down formula for 100(1-α) % confidence interval for the difference being population means (Consider independent case only) 16. A) Suppose that x₁, x₂,, x_n from a random sample from a geometric distribution parameter θ. Derive maximum likelihood estimator of θ. B) Briefly explain the general procedure for testing of hypotheses 	ameter θ. (7) ween two (3) ution with (5) (5)